

**EXTENDED ABSTRACT**  
**VI IBERIAN CONGRESS OF ICHTHYOLOGY**



**SIBIC2016**

VI CONGRESO IBÉRICO  
DE ICTIOLOGÍA / MURCIA  
21 - 24 JUNIO

# Larval habitat of sea lamprey *Petromyzon marinus* Linnaeus, 1758

Sergio Silva<sup>1\*</sup>, Sandra Barca<sup>1,2</sup>, Rufino Vieira<sup>2</sup>, M<sup>a</sup> José Servia<sup>3</sup>, David José Nachón<sup>1</sup>, Carlos Morquecho<sup>1</sup>, Lorena Lago<sup>1</sup>, Pablo Gómez<sup>1</sup>, Maitte Couto<sup>1</sup>, M<sup>a</sup> Carmen Cobo<sup>2</sup> and Fernando Cobo<sup>1,2</sup>

1. Departamento de Zooloxía, Xenética e Antropoloxía Física, Facultade de Bioloxía, Universidade de Santiago de Compostela, 15782 Santiago de Compostela, Spain
2. Estación de Hidrobioloxía “Encoro do Con”. Castroagudín-Cea. 36617 Vilagarcía de Arousa. Pontevedra, Spain
3. Departamento de Biología Animal, Biología Vegetal y Ecología, Facultad de Ciencias, Universidad de A Coruña. Campus da Zapateira s/n, 15008 A Coruña, Spain

\* correspondence to [sergio.silva@usc.es](mailto:sergio.silva@usc.es)

**Citation:** Silva S, Barca S, Vieira R, Servia MJ, Nachón DJ, Morquecho C, Lago L, Gómez P, Couto M, Cobo MC, Cobo F (2016) Larval Habitat of Sea Lamprey *Petromyzon marinus* Linnaeus, 1758. *Fishes in Mediterranean Environments* 2016.005: 3p.  
<https://doi.org/10.29094/FiSHMED.2016.005>

Lamprey population assessments are usually focused on sampling patches of larval habitat and in some cases differentiating between optimal and suboptimal habitat (Moser et al., 2007; Harvey et al., 2010). However, most studies concerning the description of larval habitat give a special attention to the granulometric analysis of the sediment, without precisely determining optimal habitat conditions for other variables (Harvey et al., 2010). Therefore, the aim of this study was to investigate the conditions (of a wide range of abiotic variables) that characterize the larval habitat of Sea lamprey *Petromyzon marinus* Linnaeus, 1758.

Surveys were carried out annually (2007-2011) in 31 locations situated in nine rivers in NW Spain: Anllóns, Eo, Lerez, Mandeo, Masma, Mera, Sar, Ulla and Umia. At each location ammocoetes were sampled using the single pass electrofishing method described by Silva et al. (2014) and anaesthetised using a solution of benzocaine (0.3 ml l<sup>-1</sup>). All individuals were measured ( $\pm 1$  mm), weighed ( $\pm 1$  g) and released again in the point of capture. A total of 48 variables, 9 hydromorphologic (at a basin scale), and 18 mesologic and 21 water quality parameters (meso and microhabitat scale), were collected to research the relation between these abiotic parameters and the larval density and biomass of *P. marinus*: river mouth orientation, river slope, basin area, annual module, basin length, elongation index, compactation index, coefficient of irregularity, sinuosity index, river width, discharge, shape index of the cross section of the river, depth of the substrate, depth over the substrate, current velocity over the

larval habitat and in the river section, distance to river mouth, vegetation cover, five granulometric fractions, mean and maximum depth of the water column in the river section, index of organic pollution, saturation rate of oxygen, dissolved oxygen, total dissolved solids, suspended solids, organic matter in the water, organic matter in the sediment, orthophosphates, nitrites, nitrates, ammonium, sulphates, turbidity, conductivity, pH, Calcium, Magnesium, hardness and chlorophyll a.

Spearman's rho ( $r_s$ ) was used to investigate the relation between abiotic variables and the density and biomass of lamprey as data did not conform to normality. Larval density and biomass were classified as: very high, high, moderate, poor or very poor following Silva et al. (2014). Abiotic characteristics were analyzed for all sites where lamprey were captured (suitable larval habitat), sites with high or very high densities or biomasses (optimal habitat) and for sites with very poor, poor or moderate densities and biomasses (suboptimal habitat). Statistical analyses were performed with IBM SPSS Statistics 20.0.

A total of 8716 lamprey were captured (8338 larvae and 378 metamorphosing individuals). Significant correlations were obtained between the larval density and biomass and the section slope ( $r_s= 0.407$ ,  $p < 0.05$  for density;  $r_s= 0.434$ ,  $p < 0.01$  for biomass) and the sinuosity index ( $r_s= 0.387$ ,  $p < 0.05$  for biomass). Significant correlations were also obtained with mesologic and physicochemical parameters (Table 1). Similar correlations were obtained for sites with high or very high larval densities or biomasses (optimal larval habitat).

Larger basins with sinuous and productive rivers hosted more abundant populations of ammocoetes, probably linked with a greater availability of food and suitable spawning and larval habitats. Areas with some organic enrichment were also favourable for ammocoetes due to the greater availability of food (Moss, 1980). However, ammocoetes were not present or were in very low densities-biomasses in areas with chronic organic pollution. The optimal habitat was constituted mainly by sand located in shallow areas (~30 cm depth) with mild organic enrichment, slow to moderate water flows (~8 cm s<sup>-1</sup>) and a good-acceptable oxygen saturation rate (>70%).

In situ differentiation between optimal and suboptimal larval habitat in population studies is not recommended, due to a large number of variables involved and the difficulty or impossibility to properly characterize them on field. It is recommended to conduct the sampling in the existing patch -or patches- of larval habitat in the selected section of river that better matches the optimal conditions.

## ACKNOWLEDGEMENTS

Facilities needed for this work were provided by the Hydrobiological Field Station "Encoro do Con" (University of Santiago de Compostela).

## CITED REFERENCES

- Harvey, J. Noble, A., Nunn, A., et al. (2010). Monitoring Sea Lamprey *Petromyzon marinus* Ammocoetes in SAC Rivers. A Case Study on the River Wye. In: Conservation Monitoring in Freshwater Habitats. Hurford, C., Schneider, M., & Cowx, I. (eds.). p. 193-206. Springer: London, UK.
- Moss, B. (1980). Ecology of freshwaters. Blackwell Scientific Publications: Oxford.
- Silva, S., Vieira-Lanero, R., Barca, S., et al. (2014). Single pass electrofishing method for assessment and monitoring of larval lamprey populations. *Limnetica*, 33 (2): 217-226.
- Moser, M. L., Butzerin, J. M., & Dey, D. B. (2007). Capture and collection of lam-
- preys: the state of the science. *Reviews in Fish Biology and Fisheries*, 17: 45-56.

**TABLE 1.** Significant correlations (Spearman's rho) obtained among physicochemical or mesologic parameters and the larval density or biomass. \*:  $p < 0.05$ ; \*\*:  $p < 0.01$ . TDS: total dissolved solids; F.: granulometric fraction.

Parameters	Density	Biomass
Mean depth	-0.304	-0.339*
pH	0.217*	0.214*
Conductivity	0.225*	0.238**
TDS	0.217*	0.231*
Hardness [CaCO <sub>3</sub> ]	0.291**	0.288**
Ca <sup>++</sup>	0.281**	0.279**
Mg <sup>++</sup>	0.244*	0.205*
Nitrates	-0.179	-0.206*
Amonium	-0.241*	-0.196*
Orthophosphates	-0.276**	-0.260**
F. [>2 mm]	0.257	0.506**
F. [1-2 mm]	0.160	0.373**
F. [0.25-0.5 mm]	-0.095	-0.331*
F. [0.125-0.25 mm]	-0.295*	-0.501**
F. [0.063-0.125 mm]	-0.283*	-0.458**
F. [<0.063 mm]	-0.253	-0.438**